

Topology and Higher-Dimensional Category Theory: the Rough Idea

Tom Leinster

Abstract

Higher-dimensional category theory is the study of n -categories, operads, braided monoidal categories, and other such exotic structures. Although it can be treated purely as an algebraic subject, it is inherently topological in nature: the higher-dimensional diagrams one draws to represent these structures can be taken quite literally as pieces of topology. Examples of this are the braids in a braided monoidal category, and the pentagon which appears in the definitions of both monoidal category and A_∞ -space.

I will try to give a Friday-afternoonish description of some of the dreams people have for higher-dimensional category theory and its interactions with topology. Grothendieck, for instance, suggested that tame topology should be the study of n -groupoids; others have hoped that an n -category of cobordisms between cobordisms between $\dots$ will provide a clean setting for TQFT; and there is convincing evidence that the whole world of n -categories is a mirror of the world of homotopy groups of spheres.

These are notes from talks given in London and Sussex in summer 2001. I thank Dicky Thomas and Roger Fenn for their invitations and the audiences for their comments, many of which are incorporated here.

What I want to give you in the next hour is an informal description of what higher-dimensional category theory is and might be, and how it is relevant to topology. There will be no real theorems, proofs or definitions. But to whet your appetite, here's a question which we'll reach an answer to by the end:

Question What is the close connection between the following two facts?


- A** No-one ever got into trouble for leaving out the brackets in a tensor product of several objects (abelian groups, etc.). For instance, it's safe to write $A \otimes B \otimes C$ instead of $(A \otimes B) \otimes C$ or $A \otimes (B \otimes C)$.
- B** There exist non-trivial knots (that is, knots which cannot be undone) in $\mathbb{R}^3$.

1 The Very Rough Idea


In ordinary category theory we have diagrams of objects and arrows such as



We can also consider more complex category-like structures, in which there are diagrams such as




This looks like an electronic circuit diagram or a flow chart; the unifying idea is that of ‘information flow’. It can be redrawn as




which looks like a surface or a diagram from topological quantum field theory.

You can also use diagrams like this to express algebraic laws, e.g. commutativity:




This is perhaps more clear when labels are added:



The fact that two-dimensional TQFTs are essentially Frobenius algebras is an example of an explicit link between the spatial and algebraic aspects of diagrams like these.

Moreover, if you allow crossings, as in the diagrams for commutativity or as in




then you start getting pictures that look like knots; and there are indeed well-established relations between knot theory and higher categorical structures.

So the idea is:


in ordinary category theory we have 1-dimensional arrows $\longrightarrow$;
in higher-dimensional category theory we have higher-dimensional arrows.

The natural geometry of these higher-dimensional arrows is what makes higher-dimensional category theory an inherently topological subject.

Example For a concrete example of what I mean by a ‘higher-dimensional arrow’, consider operads. An operad consists of operations shaped like



with many inputs (five, here) and one output. Any tree of operations, such as



can be composed to give a single operation. (Strictly speaking, these pictures are appropriate for operads without a symmetric group action: ‘non- Σ operads’, or ‘planar operads’ as they are sometimes called.)

There’s a whole zoo of structures coming under the heading of higher-dimensional category theory, including operads, generalized operads (of which the variety familiar to topologists is a basic special case), multicategories, various flavours of monoidal categories, and n -categories. Today I’ll concentrate on n -categories.

Terminology: a ‘higher-dimensional category’ or n -category is not a special kind of category, but a generalization of the notion of category; compare the usage of ‘quantum group’. A 1-category is the same thing as an ordinary category.

2 n -Categories

Here’s a very informal

‘Definition’ Let $n \geq 0$. An n -category consists of

- 0-cells or objects, $A, B, \dots$

- 1-cells or morphisms, drawn as $A \xrightarrow{f} B$

- 2-cells $A \begin{array}{c} \xrightarrow{f} \\ \Downarrow \alpha \\ \xrightarrow{g} \end{array} B$ (‘morphisms between morphisms’)

- 3-cells $A \begin{array}{c} \xrightarrow{f} \\ \alpha \circlearrowleft \Gamma \circlearrowright \beta \\ \xrightarrow{g} \end{array} B$ (where the arrow labelled Γ is meant to be going in a direction perpendicular to the plane of the paper)
- ...
- all the way up to n -cells
- various kinds of *composition*, e.g.

$$A \xrightarrow{f} B \xrightarrow{g} C \quad \text{gives} \quad A \xrightarrow{g \circ f} C \quad (\text{as we're all familiar with})$$

$$\begin{array}{ccc} \begin{array}{c} f \\ \Downarrow \alpha \\ A \xrightarrow{g} B \\ \Downarrow \beta \\ h \end{array} & \text{gives} & \begin{array}{c} f \\ \Downarrow \beta \circ \alpha \\ A \xrightarrow{h} B \end{array} \\ \begin{array}{c} f \quad f' \\ \Downarrow \alpha \quad \Downarrow \alpha' \\ A \xrightarrow{g} A' \xrightarrow{g'} A'' \end{array} & \text{gives} & \begin{array}{c} f' \circ f \\ \Downarrow \alpha' * \alpha \\ A \xrightarrow{g' \circ g} A'' \end{array} \end{array}$$


(the $*$ notation is traditional but not particularly well-chosen)

and so on in higher dimensions (which I won't attempt to draw); and similarly *identities*.

These compositions are required to 'all fit together nicely'—a phrase hiding many subtleties.

∞ -categories (also known as ω -categories) are defined similarly, by going on up the dimensions forever instead of stopping at n .

You could also consider some kind of higher categorical structure which involved cubical (or other) shapes, e.g. a 2-cell might look like



This is one of the 'zoo of structures' mentioned earlier, but is something other than an n -category.

Critical Example This is the excuse for putting the word ‘Grothendieck’ into the abstract. Any topological space X gives rise to an ∞ -category $\Pi_\infty(X)$ (its *fundamental ∞ -groupoid*), in which

- 0-cells are points of X , drawn as $\bullet$
- 1-cells are paths in X (parametrized, i.e. maps $[0, 1] \longrightarrow X$), drawn as $\bullet \longrightarrow \bullet$ —though whether that’s meant to be a picture in the space X or the ∞ -category $\Pi_\infty(X)$ is deliberately ambiguous; we’re trying to blur the distinction between geometry and algebra
- 2-cells are homotopies of paths (relative to endpoints), drawn 
- 3-cells are homotopies of homotopies of paths (i.e. suitable maps $[0, 1]^3 \longrightarrow X$)
- ...
- composition is by pasting paths and homotopies.

So $\Pi_\infty(X)$ should contain all the information you want about X if your context is ‘tame topology’; e.g. you should be able to compute from it the homotopy, homology and cohomology of X . (The word ‘groupoid’ means that all cells of dimension > 0 are invertible.)

You can also truncate after n steps in order to obtain $\Pi_n(X)$, the *fundamental n -groupoid* of X : e.g. $\Pi_1(X)$ is the familiar fundamental groupoid.

Alert As you may have noticed, composition in $\Pi_\infty(X)$ isn’t genuinely associative; nor is it unital, and nor are the cells genuinely invertible (only up to homotopy). We’re therefore interested in *weak n -categories*, where the ‘fitting together nicely’ only happens up to some kind of equivalence, rather than *strict n -categories*, where associativity etc. hold in the strict sense.

To define strict n -categories precisely turns out to be easy. To define weak n -categories, we face the same kind of challenge as algebraic topologists did in the 60s, when they were trying to state the exact sense in which a loop space is a topological group. It clearly isn’t a group in the literal sense, as composition of paths isn’t associative; but it is associative up to homotopy, and if you pick specific homotopies to do this job then these homotopies obey laws of their own—or at least, obey them up to homotopy; and so on. At least two precise formulations of ‘group up to (higher) homotopy’ became popular: Stasheff’s A_∞ -spaces and Segal’s special Δ -spaces. (More exactly, these are notions of monoid or semigroup up to homotopy; the inverses are dealt with separately.)

The situation for weak n -categories is similar but more extreme: there are something like a dozen proposed definitions that I know of, and no-one has much idea of how they relate to one another. (Some of the reasons for this chaos are good: there are real conceptual difficulties in saying what it means for two definitions of weak n -category to be equivalent.) Happily, we can ignore all this today and work informally. This means that nothing I say from now on is true with any degree of certainty or accuracy.


At this point you might be thinking: can't we do away with this difficult theory of weak n -categories and just stick to the strict ones? The answer is: if you're interested in topology, no. The difference between the weak and strict theories is genuine and nontrivial: for while it is true that every weak 2-category is equivalent to some strict one, and so it is also true that homotopy 2-types can be modelled by strict 2-groupoids, neither of these things is true in dimensions ≥ 3 . For instance, there exist spaces X such that the weak 3-category $\Pi_3(X)$ is not equivalent to any strict 3-category.

From now on, ' n -category' will mean 'weak n -category'. The strict ones hardly arise in nature.

Some More Examples of ∞ -categories:

Top This is very similar to the Π_∞ example above. **Top** has:


- 0-cells: topological spaces
- 1-cells: continuous maps



- 3-cells: homotopies between homotopies (i.e. suitable maps $[0, 1]^2 \times X \longrightarrow Y$)
- ...
- composition as expected.

ChCx This ∞ -category has:

- 0-cells: chain complexes (of abelian groups, say)
- 1-cells: chain maps
- 2-cells: chain homotopies



maps $\Gamma : A \longrightarrow B$ of degree 2 such that $d\Gamma - \Gamma d = \beta - \alpha$


- ...
- composition: more or less as expected, but some choices are involved. For instance, if you try to write down the composite of two chain homotopies $\Downarrow \Downarrow$ then you'll find that there are two equally reasonable ways of doing it: one 'left-handed', one 'right-handed'.

This is something like choosing the parametrization when deciding how to compose two loops in a space (usual choice: do everything at double speed). Somehow the fact that there's no canonical choice means that weakness of the resulting ∞ -category is inevitable.

In a reasonable world there ought to be some kind of map **Chains** : **Top** $\longrightarrow$ **ChCx**.


Bord This is an ∞ -category of (co?)bordisms.

- 0-cells: 0-manifolds, where ‘manifold’ means ‘compact, smooth, oriented manifold’. A typical 0-cell is $\bullet \bullet \bullet \bullet$.
- 1-cells: 1-manifolds with corners, i.e. cobordisms between 0-manifolds, such as



(this being a 1-cell from the 4-point 0-manifold to the 2-point 0-manifold). In Atiyah-Segal-style TQFT, we'd stop here and take *isomorphism classes* of the 1-cells just described, to make a category. We avoid this (unnatural?) quotienting out and carry on up the dimensions.

- 2-cells: 2-manifolds with corners, such as



Here N is a 2-cell $L \begin{array}{c} \xrightarrow{M} \\ \Downarrow N \\ \xrightarrow{M'} \end{array} L'$, where

$$L = L' = \bullet \bullet, \quad M = \begin{array}{c} \cup \\ \cap \end{array}, \quad M' = \begin{array}{c} | \\ | \end{array}.$$

I've left the orientations off the pictures, but N is meant to be oriented so as to agree with the orientations of M and M' . In his [2001] paper, Khovanov discusses TQFTs with corners in the language of 2-categories; he'd stop here and take isomorphism classes of the 2-cells just described, to make a 2-category. Again, we do not quotient out but keep going up the dimensions.

- 3-cells, 4-cells, ... are defined similarly
- composition is gluing of manifolds.

Some authors discuss 'extended TQFTs' via the notion of n -vector space. A 0-vector space is a complex number; a 1-vector space is an ordinary vector space; and n -vector spaces for higher n are something more sophisticated. I won't attempt to say anything about this; see Further Reading below for pointers.

3 Degenerate n -Categories

So far we've seen that topological structures provide various good examples of n -categories, and that alone might be enough to convince you that n -categories are interesting from a topological point of view. But the relationship between topology and higher-dimensional category theory is actually much more intimate than that. To see this, we'll consider n -categories which are degenerate in various ways. This doesn't sound very promising, and it'll seem at first as if it's a purely formal exercise, but in a little while the intrinsic topology should begin to shine through.

Some Degeneracies

- A category $\mathcal{C}$ with only one object is the same thing as a monoid (= semigroup with unit) M . For if the single object of $\mathcal{C}$ is called $*$, say, then $\mathcal{C}$ just consists of the set $\text{Hom}(*, *)$ together with a binary operation of composition and a unit element 1 , obeying the usual axioms. So we have:

$$\begin{aligned} \text{morphism in } \mathcal{C} &= \text{element of } M \\ \circ \text{ in } \mathcal{C} &= \cdot \text{ in } M. \end{aligned}$$



- A 2-category $\mathcal{C}$ with only one 0-cell is the same thing as a monoidal category $\mathcal{M}$. (Private thought: if $\mathcal{C}$ has only one 0-cell then there are only interesting things happening in the top two dimensions, so it's going to be *some* kind of one-dimensional structure.) This works as follows:

$$\begin{aligned}
 & \text{1-cell in } \mathcal{C} &= & \text{object of } \mathcal{M} \\
 & \text{2-cell in } \mathcal{C} &= & \text{morphism of } \mathcal{M} \\
 & \text{composition } \bullet \longrightarrow \bullet \longrightarrow \bullet \text{ in } \mathcal{C} &= & \otimes \text{ of objects in } \mathcal{M} \\
 & \text{composition } \begin{array}{c} \circlearrowleft \\ \Downarrow \\ \circlearrowright \end{array} \text{ in } \mathcal{C} &= & \circ \text{ of morphisms in } \mathcal{M}.
 \end{aligned}$$

- A monoidal category $\mathcal{C}$ with only one object is... well, if we forget the monoidal structure for a moment then we've already seen that it's a monoid whose elements are the morphisms of $\mathcal{C}$ and whose multiplication is the composition in $\mathcal{C}$. But the monoidal structure on $\mathcal{C}$ provides not only a tensor product for objects, but also a tensor product for morphisms: so the set of morphisms of $\mathcal{C}$ has a second multiplication on it, $\otimes$. So a one-object monoidal category is a set M equipped with two monoid structures which are in some sense compatible (because of the axioms on a monoidal category). And there's a well-known result (the Eckmann-Hilton argument) saying that in this situation, the two multiplications are in fact equal and commutative. So: a one-object monoidal category is a commutative monoid.

This is essentially the same argument often used to prove that the higher homotopy groups are abelian, or that the fundamental group of a topological group is abelian. In fact, we can deduce that π_2 is abelian from our 'results' so far:

Corollary: $\pi_2(X, x_0)$ is abelian (for a space X with basepoint x_0). For the 2-category $\Pi_2(X)$ has a sub-2-category whose only 0-cell is x_0 , whose only 1-cell is the constant path at x_0 , and whose 2-cells are all the possible ones from $\Pi_2(X)$ —that is, are the homotopies from the constant path to itself, that is, are the elements of $\pi_2(X, x_0)$. This sub-2-category is a 2-category with only one 0-cell and one 1-cell, i.e. a monoidal category with only one object, i.e. a commutative monoid; and this monoid is exactly $\pi_2(X, x_0)$.

- Next consider a 3-category with only one 0-cell and one 1-cell. We haven't looked at (weak) 3-categories in enough detail to work this out properly, but it turns out that such a 3-category is the same thing as a braided monoidal category. By definition, a *braided monoidal category* is a monoidal category equipped with a map (a *braiding*)


$$A \otimes B \xrightarrow{\beta_{A,B}} B \otimes A$$

for each pair A, B of objects, satisfying axioms *not* including that

$$(A \otimes B \xrightarrow{\beta_{A,B}} B \otimes A \xrightarrow{\beta_{B,A}} A \otimes B) = 1.$$


The canonical example of a braided monoidal category (in fact, the braided monoidal category freely generated by a single object) is **Braid**. This has:

- objects: natural numbers $0, 1, \dots$
- morphisms: braids, e.g.



(taken up to deformation); there are no morphisms $m \longrightarrow n$ when $m \neq n$

- tensor: placing side-by-side (which on objects means addition)
- braiding: left over right, e.g.



(Notice how $\beta_{n,m} \circ \beta_{m,n}$ is not the identity braid.)

- We're getting way out of our depth here, but nevertheless: we've already considered categories (= categories), 2-categories which are only interesting in the top two dimensions (= monoidal categories), and 3-categories which are only interesting in the top two dimensions (= braided monoidal categories). What next? For $r \geq 4$, an r -category with only one i -cell for each $i < r - 1$ is (people think) the same as a symmetric monoidal category (i.e. a braided monoidal category in which $\beta_{B,A} \circ \beta_{A,B} = 1$ for all A, B). So the situation's stabilized... and this is meant to make you start thinking of stabilization phenomena in homotopy.

The Big Picture Let's try to assemble this information on degeneracies in a systematic way. Define a k -monoidal n -category to be a $(k + n)$ -category with only one i -cell for each $i < k$. (It's clear that this is going to be some kind of n -dimensional structure, as there are only interesting cells in the top $(n + 1)$ dimensions.) Here's what k -monoidal n -categories are for some low values of k and n , laid out in the so-called 'periodic table'. Explanation follows.

		n			
		0	1	2	3
k	0	set	category	2-category	3-category
	1	monoid	monoidal category	monoidal 2-category	monoidal 3-category
	2	commutative monoid	braided mon cat	braided mon 2-cat	braided mon 3-cat
	3	“	symmetric mon cat	X	X
	4	“	“	symmetric mon 2-cat	X
	5	“	“	“	symmetric mon 3-cat
	6	“	“	“	“

↙ : take just one-object things

Running commentary: in the first row ($k = 0$), a 0-monoidal n -category is simply an n -category (it's not monoidal at all).

In the next row ($k = 1$), a 1-monoidal n -category is a monoidal n -category, i.e. an n -category equipped with a tensor product which is associative and unital up to equivalence of a suitable kind. For instance, a 1-monoidal 0-category is a one-object (1-)category, i.e. a monoid; and a 1-monoidal 1-category is a one-object 2-category, i.e. a monoidal category. We didn't look at the case of one-object 3-categories, but they turn out to be monoidal 2-categories. (What's a monoidal 2-category? Well, it's a one-object 3-category... or a direct definition can be supplied.) We see from these examples, or the general definition of k -monoidal n -category, that going in the direction ↙ means restricting to the one-object structures.

Now look at the third row ($k = 2$): we've seen that a degenerate monoidal category is a commutative monoid and a doubly-degenerate 3-category is a braided monoidal category. It's customary to keep writing 'braided monoidal n -category' all along the row, but you can regard this as nothing more than name-calling.

Next consider the first *column* ($n = 0$). A one-object braided monoidal category is going to be a commutative monoid with a little extra data (for the braiding) obeying some axioms, but it turns out that we don't actually get anything new: in some sense, 'you can't get better than a commutative monoid'. This gives the entry for $k = 3, n = 0$, and you get the same thing all the way down the rest of the column.

A similar story applies in the second column ($n = 1$). We saw—or rather, I claimed—that for $k \geq 3$, a k -monoidal 1-category is just a symmetric monoidal category. So again the column stabilizes, and again the point of stabilization is ‘the most symmetric thing possible’.

The same goes in subsequent columns. The X’s could be replaced by more terminology, e.g. the first is sometimes called ‘sylleptic monoidal 2-category’, but it doesn’t matter what that means.

The main point is that the table stabilizes for $k \geq n + 2$ —just like $\pi_{k+n}(S^k)$. So if you overlaid a table of the homotopy groups of spheres onto the table above then they’d stabilize at the same points. There are arguments to see why this should be so (and I remind you that this is all very informal and by no means completely understood). Roughly, the fact that the prototypical braided monoidal category **Braid** is not symmetric comes down to the fact that you can’t usually translate two 1-dimensional affine subspaces of 3-dimensional space past each other; and this is the same kind of dimensional calculation as you make when proving that the homotopy groups of spheres stabilize.


Answer to the initial question...

A Every weak 2-category is equivalent to a strict one. In particular, every (weak) monoidal category is equivalent to a strict one. So, for instance, we can pretend that the monoidal category of abelian groups is strict, and so that $\otimes$ is strictly associative.


B *Not* every weak 3-category is equivalent to a strict one. In fact we’ve already seen a counterexample. For

- a weak 3-category with one 0-cell and one 1-cell is a braided monoidal category
- a strict 3-category with one 0-cell and one 1-cell is a strict symmetric monoidal category
- if a braided monoidal category is equivalent to a symmetric monoidal category then it’s symmetric.

So any braided monoidal category which is not symmetric is a weak 3-category not equivalent to a strict one. An example (the canonical example?) of this is **Braid** itself; and the fact that **Braid** is not symmetric says exactly that the overpass



can’t be deformed to the underpass



in $\mathbb{R}^3$.

4 Further Reading

There are many other things written on higher-dimensional category theory. I'll only mention a few.

Another introduction to the theory, with many similar themes to this one, is the first half of Baez's [1997] paper. Baez and collaborators have written other interesting things on the interaction (still to be made rigorous, mostly) between topology and higher-dimensional category theory, and in particular his [1995] paper with Dolan contains more on TQFT, the periodic table, and stabilization.

Specifically 2-categorical approaches to TQFT can be found in Tillmann [1998] and Khovanov [2001]. n -vector spaces are to be found in Kapranov and Voevodsky [1994], and their possible role in topological field theory is discussed in Lawrence [1996].

Grothendieck puts the case that tame topology is really the study of ∞ -groupoids in his epic [1984] letter to Quillen.

One interesting idea that I didn't mention in this talk is that n th cohomology should have coefficients in an n -category (as opposed to an abelian group). This is explained in the Introduction to Street's paper of [1987].

A serious and, of course, highly recommended survey of the proposed definitions of weak n -category, including ten such definitions, is my own [2001] paper. Some other occupants of the zoo of higher-dimensional structures, including generalized operads and multicategories, are to be found in my [2000], the last chapter of my [1998], and the second half of Baez's [1997].

References

- [1997] John C. Baez, An introduction to n -categories. In *7th Conference on Category Theory and Computer Science*, ed. E. Moggi and G. Rosolini, Springer Lecture Notes in Computer Science vol. 1290 (1997). E-print [q-alg/9705009](#) .
- [1995] John C. Baez, James Dolan, Higher-dimensional algebra and topological quantum field theory. *Journal of Mathematical Physics* 36 (11) (1995), 6073–6105. E-print [q-alg/9503002](#) (without pictures).
- [1984] A. Grothendieck, Pursuing stacks (1984). Manuscript.
- [1994] M. M. Kapranov, V. A. Voevodsky, 2-categories and Zamolodchikov tetrahedra equations (1994). In *Algebraic Groups and their Generalizations: Quantum and Infinite-Dimensional Methods*, Proceedings of Symposia in Pure Mathematics 56, Part 2, AMS, 177–260.
- [2001] Mikhail Khovanov, A functor-valued invariant of tangles (2001). E-print [math.QA/0103190](#) .
- [1996] R. J. Lawrence, An introduction to topological field theory (1996). In *The Interface of Knots and Physics*, Proceedings of Symposia in Applied Mathematics 51, AMS, 89–128. Also available via <http://www.ma.huji.ac.il/~ruthel> .

- [1998] Tom Leinster, Structures in higher-dimensional category theory (1998). Available via <http://www.dpmms.cam.ac.uk/~leinster> .
- [2000] Tom Leinster, Operads in higher-dimensional category theory. PhD thesis, University of Cambridge, 2000. E-print [math.CT/0011106](#) .
- [2001] Tom Leinster, A survey of definitions of n -category (2001). E-print [math.CT/01xxxxx](#) (to appear shortly).
- [1987] Ross Street, The algebra of oriented simplexes. *Journal of Pure and Applied Algebra* 49 (1987), no. 3, 283–335.
- [1998] Ulrike Tillmann, $\mathcal{S}$ -structures for k -linear categories and the definition of a modular functor. *Journal of the London Mathematical Society (2)* 58 (1998), no. 1, 208–228. E-print [math.GT/9802089](#) .

Department of Pure Mathematics, University of Cambridge

Email: leinster@dpmms.cam.ac.uk

Web: <http://www.dpmms.cam.ac.uk/~leinster>